MBBS (HONS), BBIOMEDSC (HONS), FRACP | GASTROENTEROLOGIST
ARGUS DRJEREMYDWYER@ARGUS.NET.AU | HEALTHLINK DRJDWYER

EXAMINATION REQUESTED ☐ Gastroscopy ☐ Colonoscopy ☐ Consultation ☐ Other **PATIENT DETAILS** ☐ Mrs Ms Given Name Surname D.O.B Address Postcode Phone Mobile Email REFERRING DOCTOR Affix stamp here or include name, address, provider number

Consulting locations:

Suite 5.3 Level 5, 124 Grey Street (Epworth Freemasons Hospital) East Melbourne VIC 3002

• 438 High Street, Melton VIC 3337

(All correspondence to): PO Box 68, Yarraville VIC 3013

P 03 9996 9545 | **M** 0435 402 789 | **F** 03 9492 5282 | **E** reception@drjeremydwyer.com

CLINICAL NOTES	
Cianad	Date
Signed	Date
ANTI-COAGULANT/ANTI-PL	ATELET
□ No □ Yes	Rx:
DIABETES	
□ No □ Yes	Rx:
MORBID OBESITY BMI >40	
□ No □ Yes	Weight:
ALLERGIES	
□ No □ Yes	

GENERAL INSTRUCTIONS FOR GASTROSCOPY/COLONOSCOPY:

Please bring this referral or your doctor's referral on the day of your procedure and any private health insurance details, Medicare card, DVA card and medication list. For patients undergoing colonoscopy, you will be provided with specific instructions for the bowel preparation, which should be purchased from your local pharmacist at least 3 days prior to your procedure. On the day of your procedure, you must have nothing to eat or drink for at least six hours beforehand apart from 'clear liquids' which must be ceased at least two hours prior to your procedure. Any important regular medications can be taken with a sip of water on the day of the procedure but do not take any diabetic medications. You will be admitted for approximately 3 hours for your procedure. Because of the sedation medication given during the procedure, it is very important that you do not drive a car, travel on public transport or catch a taxi/uber alone, operate machinery, sign legal documents or drink alcohol within the first 24 hours after the procedure. It is mandatory that a responsible adult accompanies you home after the procedure and stays with you overnight. Although complications can occur with any procedure, they are very uncommon following diagnostic gastroscopy and colonoscopy (approximately 1 in 1000 cases).